

NEAR SURFACE EMPLACEMENT OF KIMBERLITES: CONTRASTING MODELS AND WHY

Field, M¹ and Scott Smith, B.H.²

1. De Beers Consolidated Mines, Kimberley Mines, P.O. Box 10191, Beaconsfield, 8315, South Africa.
2. Scott Smith Petrology Inc., 2555 Edgemont Boulevard, North Vancouver, B.C. V7R 2M9., Canada

The most complete kimberlite emplacement models result from detailed studies of specific localities. Investigations undertaken for economic reasons create extensive 3D exposure and permit thorough studies. The first such model outside Russia was developed for the main diamond mining areas of that time around Kimberley, South Africa (Clement and Reid, 1989 and previous work by Clement and Skinner). These studies showed that kimberlites are unusual, and require kimberlite-specific terminology (Field and Scott Smith, this volume), and emplacement models. The Cretaceous southern African kimberlite pipes were shown to comprise three zones: a complex root zone, a steep sided diatreme zone and an overlying crater zone which together can be up to 2km deep. Each zone is largely filled with texturally specific types of kimberlite (Fig. 2 of Field and Scott Smith, this volume), indicating that each zoned formed by a different process.

The model for these kimberlites shows that the near surface emplacement occurred in a closed system created by many barriers and cap rocks through which the kimberlite has to intrude (e.g. Karoo dolerites, basalts, earlier kimberlites). The kimberlite magma was clearly inhibited during its final ascent to surface resulted in complex and massive subsurface brecciation of the country rock as well as an immense build up of juvenile gases, volumetrically dominated by carbon dioxide. When breakthrough to surface was achieved, sudden and rapid degassing caused fluidization of the subsurface magma. This in turn excavated the diatreme zone through the pre-brecciated country rock. This process was short lived and the fluidized, degassed magma rapidly condensed to form a very specific textural rock type (tuffitic kimberlite breccia). This rock type contains in the order of 50% xenoliths derived from the adjacent pipe wall which are set in the products of the remaining condensed kimberlitic fluids. It is important to emphasize that the same process created both the diatreme and its infilling. The underlying root zone and overlying crater zone formed by more normal processes that are part of the same event. This model has been tested and validated by other workers. At Orapa, Botswana the same model can be extended up into the crater (Field et al., 1997). At the Orapa North pipe a single extremely powerful magmatic eruption caused the excavation of both the diatreme and crater. The crater was rapidly infilled with primary pyroclastic products, that are similar to, and derived from, the fluidized kimberlite that formed the diatreme below. It has been shown elsewhere in the world that this model has been repeated in space and time.

In contrast Lorenz (1985 and other papers) proposed a phreatomagmatic "maar-diatreme" model for the southern African kimberlite pipes. Maars are formed by two separate processes : (1) crater excavation and (2) subsequent infilling. The maar craters are excavated into the country rock by phreatomagmatic explosions that relate to an aquifer. The products of this process form mainly crater rim base surge type deposits. Subsequent infilling of the crater is mainly by re-sedimentation of crater rim or wall material, although there may be some infill derived from pyroclastic magmatic eruptions. The latter notably lack country rock xenoliths. Many recent maars are only partially infilled. The same maar-forming process is repeated in space and time. Notably, in many cases there seems to be scant evidence for the presence of steep sided pipes below maars. There is no geological

evidence to support the suggestion that maar-like phreatomagmatic processes were important in the formation and infilling of the southern African kimberlite pipes.

Scott-Smith et al. (this volume) have developed a model for the Cretaceous kimberlites in Saskatchewan, Canada, which differs considerably from that of the southern African kimberlite diatreme model. No diatremes or root zones were developed in these pipes. The pipes formed by two separate processes : (1) crater excavation and (2) subsequent infilling. Empirically these saucer shaped craters are thought to have formed by maar-like phreatomagmatic processes relating to a well-known aquifer approximately 200m below the present surface, the point from which the craters flare. It is considered that the lack of barriers and cap rocks in the 600m of Phanerozoic sedimentary cover permitted the kimberlite magmas to reach surface easily. The kimberlite magma, therefore, erupted in an open system. This precluded the sub-surface build up of juvenile volatiles which is requisite for the formation of diatreme and root zones. Subsequent to crater formation, the craters were filled with primary pyroclastic kimberlite and re-sedimentation does not appear to have been an important process. In this respect, the Saskatchewan pipes differs from maars. In Saskatchewan much of the infill was derived from Hawaiian and Strombolian-style eruptions. In a few pipes a different kimberlite-specific eruption mechanism has been proposed. This more explosive style of eruption resulted from the massive degassing of juvenile volatiles close to surface and formed mega-graded beds up to at least 90m thick within the craters. This and other features highlight the fact that kimberlite magmas are very different from more common magma types.

Other examples of the Saskatchewan model may occur in Zaire (Scott Smith et al., this volume) and in Alberta at Mountain Lake (Wood et al., this volume) and Buffalo Hills (Carlson et al., this volume). In this area of Alberta 1-2km of Phanerozoic sediments overlie the basement. No diatreme-facies kimberlite has been identified at these localities but the Cretaceous pipes are infilled with xenolith-poor volcanoclastic kimberlite. It is therefore presumed that these pipes were excavated by maar-like processes. In contrast to Saskatchewan, some of the Alberta pipes appear to be steeper sided and/or deeper (up to 350m). The different shape may reflect the variable nature of the sediments. In Saskatchewan the saucer-like shapes reflect the shallow location of the aquifer as well as the poorly consolidated and fine grain size of the uppermost Cretaceous sediments. In Alberta the uppermost Cretaceous sediments are sandier and so can support steeper crater walls. The fact that these pipes penetrate further than in Saskatchewan implies that a deeper aquifer must have been involved in their formation. These localities together appear to comprise a Canadian Prairies (+/- Zaire) kimberlite emplacement model where thick piles of poorly consolidated sediments containing no barrier-forming igneous rocks offer an easy route to surface for kimberlite magmas.

Recently many new kimberlites have been discovered on the Slave craton in the NWT, Canada. These kimberlites can be grouped in terms of age and geographic distribution. The youngest group (Cretaceous to Tertiary) occur in the Lac de Gras area of the central Slave. Many of these kimberlite pipes appear to be steep sided bodies with shapes superficially similar to the southern African diatremes. Relative to the southern African pipes, those at Lac de Gras appear to be small and infilled mainly with xenolith-poor volcanoclastic kimberlite (primary pyroclastic and resedimented). These Lac de Gras pipes appear to have formed by two process : pipe excavation and pipe infilling. It seems unlikely that such deep pipes (up to +/-600m) represent explosion craters formed within the Archaean basement, irrespective of the process. The similarity in pipe shape to the southern African diatremes makes it tempting to suggest that a similar fluidisation process occurred. However, no comparable diatreme-facies infill (tuffisitic kimberlite breccia) is present. It seems to be significant that the Lac de Gras pipes are at least an order of magnitude smaller than many of those in southern

Africa. This suggests that, if the Lac de Gras pipes formed by the same processes as the southern African diatremes, they resulted from less powerful eruptions. The Archaean basement could impose a difficult route to surface for the kimberlite magma. This would allow sub-surface build up of juvenile volatiles that possibly could form a diatreme. The country rock at the time of emplacement included poorly consolidated clay-rich sediments (probably <100m) directly overlying the Archaean basement. The cover sediments would have allowed an easy breakthrough to surface and may even have contained an aquifer to aid crater formation. This contrasts with the southern African Cretaceous situation where the uppermost country rocks at the time of emplacement were competent basalts lavas. These basalts formed a major final barrier for the kimberlite magmas to penetrate. The lack of such a cap rock at Lac de Gras would have limited the amount of sub-surface volatile build up and would result in the formation of overall smaller diatremes. Some pipes in the Lac de Gras area contain only hypabyssal kimberlite. It is interesting to speculate why these kimberlite magmas did not excavate pipes similar to the volcanoclastic-infilled bodies.

Elsewhere in the Slave the known kimberlites appear are older (~172-538 Ma.) and so pre-date the Cretaceous sediments that formed the cover in the Lac de Gras area. In contrast it appears that in the other areas of the Slave the cover sediments were dominated by Palaeozoic carbonates. Although volcanoclastic and hypabyssal kimberlites are also present, there has also been development of southern African-style diatremes infilled with the diagnostic tuffisitic kimberlite breccias.

Conclusions : Kimberlites are unique carbon dioxide-rich magmas, whose near surface emplacement is controlled by the abundance juvenile volatiles and country rock geology. Two extreme emplacement models have been identified: (1) the southern African model and (2) the Canadian Prairies model. In some near surface environments, such as in southern African, barrier and cap rocks retard the upward movement of the magma. This produces closed magmatic systems from which kimberlite-specific magmatic processes generate pipes with distinctive root, diatreme and crater zones, each filled with different textural variants of kimberlite. In contrast, where the country rocks are easily penetrated by the upward moving magma, such as in the Prairies, the resulting open system precludes the build up of juvenile volatiles. Phreatomagmatic processes dominate to form maar-like craters with no diatreme or root zones. The shape of the crater depends on the nature of the country rocks and the depth of the aquifer. The infilling can be either primary pyroclastic deposits with their own kimberlite-specific characteristics and/or re-sedimented volcanoclastics. Obviously intermediate types of pipes must result from combinations of these two end member processes and the newly discovered pipes of the Lac de Gras area of the Slave may be such an example.

References

- Clement, C.R. and Reid, A.M., 1989, The origin of kimberlite pipes: an interpretation based on a synthesis of geological features, displayed by southern African occurrences. In *Kimberlites and Related Rocks*. Edited by J. Ross. Geol. Soc. of Australia Special Publication, 14, 1, p. 632-646.
- Field, M., Gibson, J.G., Wilkes, T.S., Gababitse, J. & Khujwe, P., 1995, The geology of the Orapa A/K1 kimberlite, Botswana: further insight into the emplacement of kimberlite pipes. Extended Abstracts from the 6th International Kimberlite Conference., Novosibirsk, Russia. Published by Siberian Branch of Russian Academy of Sciences, 155-157.
- Lorenz, V., 1985, Maars and diatremes of phreatomagmatic origin, *Trans, Geol. Soc., South Africa*, 88, 459-470.

SEVENTH INTERNATIONAL KIMBERLITE CONFERENCE

Cape Town, April 1998

EXTENDED ABSTRACTS

All extended abstracts in this volume were submitted by the authors in camera ready form. There has been no editing of the extended abstracts and they have not been subject to review. The Organising Committee of the 7th International Kimberlite Conference takes no responsibility for authors' errors and omissions.

SOUTH AFRICA

Author	Title	Page No's
Abdrakhimov, M.Z., Kouznetsova, E.I., Traskin, V.Yu.	Development of Intergranular cracking and relaxation of stress in deep crystalline rock of earth crust as result of physico-chemical influence of water.	1
Abe, N., Aral, S., Yurimoto, H.	Petrology of the arc peridotite xenoliths: Implications for geochemical evolution of the wedge mantle.	4
Afanasiev, V.P., Pokhilenko, N.P., Logvinova, A.M., Yefimova, E.S.	Problem of false indicators for kimberlites and lamproïtes (on the example of chromites).	7
Agashev, A.M., Fomin, A.S., Watanabe, T., Pokhilenko, N.P.	Preliminary age determination of recently discovered kimberlites of the Siberian kimberlite province.	9
Agashev, A.M., Watanabe, T., Kullgin, S.S., Pokhilenko, N.P., Orhashi, Y.	Sr-Nd isotopes in the garnet-pyroxenite xenoliths from Siberian kimberlites: a new insight into lithospheric mantle.	11
Ananiev, V.A., Kullgin, S.S., Reimers, L.F., Khlestov, V.V.	Paragenetic analysis of the upper mantle minerals from the heavy mineral concentrates of kimberlites based on an original statistic method.	14
Andre, L., Shatsky, V.S., De Corte, K., Sobolev, N.V., Navez, J., Jagoutz, E.	K - rich clinopyroxenes as mantle conveyers of crustal-derived components	17
Andronikov, A.V., Foley, S.F., Meizer, S.	Mantle xenoliths from the Jetty Peninsula area (East Antarctica): samples of thermally eroding lithosphere on the flanks of the Lambert-Amery Rift.	20
Antonyuk, B.P., Mironov, V.P.	Three-Stage Growth Model of the Natural Diamond of Octahedral Habit.	23
Araujo, A.L.N., Gaspar, J.C., Bizzi, L.C.	Petrography and mineralogy of kimberlites and kamafugites from the Alto Paranaiba Igneous Province, Minas Gerais, Brazil: Preliminary studies.	26
Araujo, D.P., Gaspar, J.C., Garg, V.K.	The Complete Phlogopite-Tetraferriphlogopite Series in the Catalao-I and -II Carbonatite Complexes, Brazil.	29
Arima, M.	Experimental Study of Growth and Resorption of Diamond in Kimberlitic Melts at High Pressures and Temperatures.	32
Ashchepkov, I., Salters, V., Andre, L.	Relationships between garnet and clinopyroxene in Vitim mantle xenoliths: evidence of the polystage growth and melt percolation?	35
Bailey, L., Helmstaedt, H.H., Peterson, R.C., Mandarino, J.A., Letendre, J.P.	Microdiamonds and Indicator Minerals from a Talc Schist Host Rock, French Guiana.	37
Barashkov, Y.P., Talnikova, S.B.	The features of the morphology crystalline inclusions and the real structure of diamond.	40
Barron, K.M., Logvinova, A.M., Sobolev, N.V.	Morphology and Composition of Chromite Macrocrysts and their Inclusions, Guaniamo Kimberlite Field, Venezuela.	43
Barron, L.M., Lishmund, S.R., Oakes, G.M., Barron, B.J.	Primary and related diamond occurrences within a Phanerozoic subduction regime in eastern New South Wales, Australia.	46
Bary, T.L., Kempton, P.D., Saunders, A.D., Windley, B.	Mantle Dynamics beneath Mongolia: Implications from Cenozoic and Mesozoic Alkalic Basalts.	49
Barth, M.G., Rudnick, R.L., Spicuzza, M.J., Valley, J.W., Haggerty, S.E.	The role of eclogites in the growth of Archean cratons: a case study from west Africa.	52
Baumgartner, M.C., Neuhoﬀ, L.	The vertical distribution of indicator minerals within Kalahari cover overlying a kimberlite pipe.	55
Beard, A.D., Mason, P.R.D., Downes, H.	Depletion and enrichment processes in lithospheric mantle beneath the Baltic Shield (Kola peninsula and Arkhangelsk) - evidence from laser ICP-MS analyses of mantle minerals.	58
Beard, A.D., Milledge, H.J.	Infrared & Microprobe studies of intrusions & micro-inclusions in diamond.	61
Bell, D.R., Mofokeng, S.W.	Cr-poor Megacrysts from the Frank Smith Mine and the source regions of transitional kimberlites	64
Belousova, E.A., Griffin, W.L., O'Reilly, S.Y.	Trace element composition and cathodoluminescence properties of kimberlitic zircons.	67
Ben Ismail, W., Mainprice, D.	An olivine fabric database: an interpretative frame for upper mantle seismology.	70
Ben Ismail, W., Mainprice, D., Barruol, G., Boyd, J., Vauchez, A.	Lithospheric Mantle anisotropy of the Kaapvaal craton (South Africa) from lattice preferred orientation analysis.	73
Berg, G.W.	Geochemical Relations which reflect the History of Kimberlites from the Type area of Kimberley, South Africa	76
Berg, G.W.	Mineralogical Evidence for, and Experimental Evidence against, Carbonate from the mantle in Xenoliths from Kimberlite Pipes.	79
Berg, G.W., Carlson, J.A.	The Leslie Kimberlite Pipe of Lac de Gras, Northwest Territories, Canada: Evidence for Near Surface Hypabyssal Emplacement.	81
Berryman, A.K., Stiefenhofer, J., Shee, S.R., Wyatt, B.A.	The Discovery and Geology of the Timber Creek Kimberlites, Northern Territory, Australia.	84
Bizzi, L.A., Pimentel, M.	Source Characteristics of Brazilian Kimberlites.	87
Bordon, V.	Mineralogical and Chemical Model of Belarussian Diatremes.	89
Bordon, V., Astapenko, V.	Braslav Field: the prospects for discovering diamondiferous rocks.	90
Bovkun, A.V., Garanin, V.K., Kudriavtseva, G.P., Possukhova, T.V.	Chemical-genetic classification of microcrystalline oxides from kimberlite groundmass and new system of diamondiferous prospecting	91
Bovkun, A.V., Garanin, V.K., Kudriavtseva, G.P., Possukhova, T.V.	Chemical-genetic classifications of oxides from kimberlite groundmass as a basis of a new technology of evaluation of diamond potential of kimberlite bodies (on an example of Yakutian and Arkhangelsk diamondiferous provinces).	94
Bovkun, A.V., Garanin, V.K., Kudriavtseva, G.P., Possukhova, T.V.	Diamonds from Timan placers: morphology, spectroscopy and genesis.	97
Boyd, F.R., Pearson, D.G., Mertzman, S.A.	Spinel-facies Peridotites from the Kaapvaal Root.	100

Author	Title	Page No's
Brown, J.W., Butcher, A.R.	Textural and Petrological Variation Within the Crater Facies Kimberlite Bodies of the Fort à la Come Province, Saskatchewan, Canada.	103
Brown, R.W., Gallagher, K., Griffin, W.L., Ryan, C.G., de Wit, M.C.J., Belton, D.X., Harman, R.	Kimberlites, accelerated erosion and evolution of the lithospheric mantle beneath the Kaapvaal craton during the mid-Cretaceous.	105
Budaev, D.A.	<<Populational>> model of kimberlites: an application to diamondiferous kimberlites of regions with various geodynamic history.	108
Budaev, D.A., Dolgunin, A.V. Fomin, A.S.	An algorithm of kimberlite diamondiferousness estimations.	111
Bulanova, G.P., Griffin, W.L., Kaminsky, F.V., Davies, R., Ryan, C.R., Andrew, A., Spetsius, Z.V., Zharkarchenko, O.D.	Diamonds from Zamitsa and Dainaya kimberlites (Yakutia): Their nature, growth history, and lithospheric mantle source.	113
Bulanova, G.P., Shelkov, D.	Nature of eclogitic diamonds from Yakutian kimberlites: evidence from isotopic composition and sulphide inclusion chemistry.	116
Burgess, R., Phillips, D., Harris, J.W., Robinson, D.N.	Antarctic diamonds in South-Eastern Australia? Hints from ⁴⁰ Ar/ ³⁹ Ar Laser probe dating of clinopyroxene inclusions from Copeton diamonds.	119
Burgess, S.R., Harte, B.	Tracing Lithosphere Evolution through the analysis of Heterogeneous G9/G10 Garnets in Peridotite Xenoliths.	122
Carlson, J.A. Kirkley, M.B., Thomas, E.M. Hillier, W.D.	Recent major kimberlite discoveries in Canada.	127
Carlson, R.W., Irving, A.J., Hearn, B.C.Jr.	Peridotite Xenoliths from the Williams Kimberlite, Montana: Implications for Delamination of the Wyoming Craton Lithosphere	132
Carlson, R.W., Pearson, D.G., Boyd, F.R., Shirey, S.B., Irvine, G., Menzies, A.H., Gurney, J.J.	Regional Age Variation of the Southern African Mantle: Significance for Models of Lithospheric Mantle Formation	135
Carlson, S.M., Hillier, W.D., Hood, C.T., Pryde, R.P., Skelton, D.N.	The Buffalo Hills Kimberlite Province, North-central Alberta, Canada.	138
Cartigny, P., Harris, J.W., Javoy, M.	Eclogitic, Peridotitic, Metamorphic Diamonds and the Problems of Carbon Recycling.	141
Channer, D.M.deR., Cooper, R.E.C., Kaminsky, F.	The Guaniamo diamond region, Bolivar state, Venezuela: a new kimberlite province.	144
Cherny, S.D., Fomin, A.S., Yanygin, J.T., Banzeruk, V.I., Kormilova, V.P.	Geology and composition of the Nakyn field kimberlite pipes and diamond properties (Yakutia).	147
Chestey, J.T., Rudnick, R.L., Lee, C.-T.	Longevity of Cratonic Mantle Beneath an Active Rift: Re-Os evidence from Xenoliths from the Tanzanian East African Rift.	149
Chinn, I.L., Gurney, J.J., Harte, B., Fitzsimmons, I.C.W., Milledge, H.J.	Nitrogen contents of diamond plates: a comparison of FTIR and SIMS analyses	152
Chinn, I.L., McCallum M.E., Harris C., Milledge, H.J., Gurney, J.J.	C02 - bearing diamonds in eclogite xenoliths from the Sloan 2 kimberlite, Colorado	155
Chinn, I.L., Milledge, H.J., Gurney, J.J.	Diamonds and inclusions from Jagersfontein kimberlite	156
Clarke, J., Sobie, P.A., Wilkes, T.A., Zweistra, P.	The Geology and Economic evaluations of the Liqhobong kimberlites, Lesotho.	158
Cookerbo, H.O.,	Emplacement history of the Jericho kimberlite pipe, northern Canada.	161
Cookerbo, H.O., Kopylova, M.G., Daoud, D.K.	A chemically and texturally distinct layer of diamondiferous eclogite beneath the central Slave craton, northern Canada.	164
Corbett, I.B., McMillan, I.K.	From Shore to Shelf and Back Again.	167
Davies, R., Griffin, W.L., Pearson, N.J., Andrew, A., Doyle, B.J., O'Reilly, S.Y.	Diamonds from the Deep: Pipe DO-27, Slave Craton, Canada.	170
Davies, R.M., O'Reilly, S.Y., Griffin, W.L.	Characteristics of Alluvial Diamonds from Bingara and Wellington, Eastern Australia.	173
Davies, R.M., O'Reilly, S.Y., Griffin, W.L.	Dynamic Growth Structures in Diamonds from Bingara, NSW, Australia.	176
Dawson, J.B.	Melting and Metasomatism in Upper Mantle Peridotite Xenoliths from Labait, North-central Tanzania, and contrasting Metasomatic styles in the Tanzanian Lithospheric Mantle.	179
de Bruin, D.	Inclusion Bearing Cr-poor and Cr-rich Garnet Megacrysts from the Group II Swartruggens Kimberlite.	181
De Corte, K., Cartigny, P., Shatsky, V.S., De Paepe, P., Sobolev, N.V., Javoy, M.	Microdiamonds from UHP Metamorphic rocks of the Kokchetav Massif, Northern Kazakhstan: FTIR spectroscopy, C & N Isotopes and Morphology.	184
Deines, P.	Intra- and Inter-Mineral Oxygen Isotope Variations in Kimberlitic Zircon.	187
Demaiffe, D., El Fadili, S., Andre, L.	Geochemical and isotopic (Sr, Nd) study of eclogite nodules from the Mbuji Mayi kimberlite, Kasai, Congo. Nature of the protoliths and evidence for mantle metasomatism.	190
De Meillon, L., Bristow, J.W.	Some Characteristics of High Level Tertiary Age Alluvial Terraces along the Orange River between the towns of Douglas and Prieska, Northern Cape Province, South Africa.	193
De Wit, M.C.J., Morelli, C., Skinner, C.P.	A reinterpretation of the Lichtenburg diamond deposits.	195
Dmitriev, A.N., Dyatlov, V.L., Litasov, K.D.	Physical model of kimberlite pipes formation: new constraints from theory of non-homogenous physical vacuum.	196
Doyle, B.J., Kivi, K., Scott Smith, B.H.	The Tili Kwi Cho (DO27 and DO18) Diamondiferous Kimberlite Complex Slave Craton, Northwest Territories, Canada	199
Elder, E., Winter, F., Edwards, R.	The Rosario do Sul Kimberlitic Province, Rio Grande do Sul State, Southern Brazil.	202
El Fadili, S., Demaiffe, D.	Petrology, mineral chemistry and thermobarometry of eclogite nodules from the Mbuji Mayi kimberlite, Kasai, Congo: significance of kyanite-cpx intergrowths.	205

Author	Title	Page No's
Brown, J.W., Butcher, A.R.	Textural and Petrological Variation Within the Crater Facies Kimberlite Bodies of the Fort à la Corne Province, Saskatchewan, Canada.	103
Brown, R.W., Gallagher, K., Griffin, W.L., Ryan, C.G., de Wit, M.C.J., Belton, D.X., Harman, R.	Kimberlites, accelerated erosion and evolution of the lithospheric mantle beneath the Kaapvaal craton during the mid-Cretaceous.	105
Budaev, D.A.	<<Populational>> model of kimberlites: an application to diamondiferous kimberlites of regions with various geodynamic history.	108
Budaev, D.A., Dolgunin, A.V. Fomin, A.S.	An algorithm of kimberlite diamondiferousness estimations.	111
Bulanova, G.P., Griffin, W.L., Kaminsky, F.V., Davies, R., Ryan, C.R., Andrew, A., Spetsius, Z.V., Zakharchenko, O.D.	Diamonds from Zamitsa and Dafnaya kimberlites (Yakutia): Their nature, growth history, and lithospheric mantle source.	113
Bulanova, G.P., Shelkov, D.	Nature of eclogitic diamonds from Yakutian kimberlites: evidence from isotopic composition and sulphide inclusion chemistry.	116
Burgess, R., Phillips, D., Harris, J.W., Robinson, D.N	Antarctic diamonds in South-Eastern Australia? Hints from ⁴⁰ Ar/ ³⁹ Ar Laser probe dating of clinopyroxene inclusions from Copeton diamonds.	119
Burgess, S.R., Harte, B	Tracing Lithosphere Evolution through the analysis of Heterogeneous G9/G10 Garnets in Peridotite Xenoliths.	122
Carlson, J.A. Kirkley, M.B., Thomas, E.M. Hillier, W.D.	Recent major kimberlite discoveries in Canada.	127
Carlson, R.W., Irving, A.J., Hearn, B.C.Jr.	Peridotite Xenoliths from the Williams Kimberlite, Montana: Implications for Delamination of the Wyoming Craton Lithosphere	132
Carlson, R.W., Pearson, D.G., Boyd, F.R., Shirey, S.B., Irvine, G., Menzies, A.H., Gurney, J.J.	Regional Age Variation of the Southern African Mantle: Significance for Models of Lithospheric Mantle Formation	135
Carlson, S.M., Hillier, W.D., Hood, C.T., Pryde, R.P., Skelton, D.N	The Buffalo Hills Kimberlite Province, North-central Alberta, Canada.	138
Cartigny, P., Harris, J.W., Javoy, M	Eclogitic, Peridotitic, Metamorphic Diamonds and the Problems of Carbon Recycling.	141
Channer, D.M.deR., Cooper, R.E.C., Kaminsky, F.	The Guaniamo diamond region, Bolivar state, Venezuela: a new kimberlite province.	144
Cherny, S.D., Fomin, A.S., Yanygin, J.T., Banzeruk, V.I., Kornilova, V.P.	Geology and composition of the Nakyn field kimberlite pipes and diamond properties (Yakutia).	147
Chesley, J.T., Rudnick, R.L., Lee, C.-T.	Longevity of Cratonic Mantle Beneath an Active Rift: Re-Os evidence from Xenoliths from the Tanzanian East African Rift.	149
Chinn, I.L., Gurney, J.J., Harte, B., Fitzsimmons, I.C.W., Milledge, H.J.	Nitrogen contents of diamond plates: a comparison of FTIR and SIMS analyses	152
Chinn, I.L., McCallum M.E., Harris C., Milledge, H.J., Gurney, J.J.	CO ₂ - bearing diamonds in eclogite xenoliths from the Sloan 2 kimberlite, Colorado	155
Chinn, I.L., Milledge, H.J., Gurney, J.J.	Diamonds and inclusions from Jagersfontein kimberlite	156
Clarke, J., Sobie, P.A., Wilkes, T.A., Zweistra, P	The Geology and Economic evaluations of the Lihobong kimberlites, Lesotho.	158
Cookenboo, H.O.,	Emplacement history of the Jericho kimberlite pipe, northern Canada.	161
Cookenboo, H.O., Kopylova, M.G., Daoud, D.K.	A chemically and texturally distinct layer of diamondiferous eclogite beneath the central Slave craton, northern Canada.	164
Corbett, I.B., McMillan, I.K	From Shore to Shelf and Back Again.	167
Davies, R., Griffin, W.L., Pearson, N.J., Andrew, A., Doyle, B.J., O'Reilly, S.Y.	Diamonds from the Deep: Pipe DO-27, Slave Craton, Canada.	170
Davies, R.M., O'Reilly, S.Y., Griffin, W.L.	Characteristics of Alluvial Diamonds from Bingara and Wellington, Eastern Australia.	173
Davies, R.M., O'Reilly, S.Y., Griffin, W.L.	Dynamic Growth Structures in Diamonds from Bingara, NSW, Australia.	176
Dawson, J.B.	Melting and Metasomatism in Upper Mantle Peridotite Xenoliths from Labait, North-central Tanzania, and contrasting Metasomatic styles in the Tanzanian Lithospheric Mantle.	179
de Bruin, D	Inclusion Bearing Cr-poor and Cr-rich Garnet Megacrysts from the Group II Swartuggens Kimberlite.	181
De Corte, K., Cartigny, P., Shatsky, V.S., De Paepe, P., Sobolev, N.V., Javoy, M.	Microdiamonds from UHP Metamorphic rocks of the Kokchetav Massif, Northern Kazakhstan: FTIR spectroscopy, C & N Isotopes and Morphology.	184
Deines, P	Intra- and Inter-Mineral Oxygen Isotope Variations in Kimberlitic Zircons.	187
Demaiffe, D., El Fadili, S., Andre, L.	Geochemical and isotopic (Sr, Nd) study of eclogite nodules from the Mbuji Mayi kimberlite, Kasai, Congo. Nature of the protoliths and evidence for mantle metasomatism.	190
De Meillon, L., Bristow, J.W	Some Characteristics of High Level Tertiary Age Alluvial Terraces along the Orange River between the towns of Douglas and Prieska, Northern Cape Province, South Africa.	193
De Wit, M.C.J., Morelli, C., Skinner, C.P.	A reinterpretation of the Lichtenburg diamond deposits.	195
Dmitriev, A.N., Dyatlov, V.L., Litasov, K.D.	Physical model of kimberlite pipes formation: new constraints from theory of non-homogenous physical vacuum.	196
Doyle, B.J., Kivi, K., Scott Smith, B.H.	The Til Kwi Cho (DO27 and DO18) Diamondiferous Kimberlite Complex Slave Craton, Northwest Territories, Canada	199
Elder, E., Winter, F., Edwards, R.	The Rosario do Sul Kimberlitic Province, Rio Grande do Sul State, Southern Brazil.	202
El Fadili, S., Demaiffe, D.	Petrology, mineral chemistry and thermobarometry of eclogite nodules from the Mbuji Mayi kimberlite, Kasai, Congo: significance of kyanite-cpx intergrowths.	205

Author	Title	Page No's
Ertinçek, Y.M., Milshtein, E.D., Saltykov, O.G., Verzhak, V.V.	Local depressions in Country Rock of Kimberlites as a New Exploration Criteria (by the Example of Zolotitsa Field, Arkhangelsk, Russia).	208
Field, M., Scott Smith, B.H.	Near surface emplacement of kimberlites: contrasting models and why.	211
Field, M., Scott Smith, B.H.	Textural and Genetic Classification schemes for kimberlites: a new perspective	214
Foley, S.F., Glaser, S.M., Andronikov, A.V.	Non-cratonic garnet peridotites from rifted continental settings in Vitim, Siberia (Baikal Rift and East Antarctica) (Lambert-Amery Rift).	217
Foley, S.F., Musselwhite, D.S., van der Laan, S.R.	Melting processes in veined lithospheric mantle in cratonic and non-cratonic settings.	220
Friese, A.E.W.	Structural control on kimberlite genesis and crustal emplacement within South Africa and the Kaapvaal Craton during the Cretaceous	224
Friese, A.E.W.	Tectonic evolution and intra-cratonic alkaline magmatism within the central Kaapvaal Craton during the Mesoproterozoic.	227
Fung, A.T.	Petrochemistry of upper mantle eclogites from the Grizzly, Leslie, Pigeon and Sable kimberlites in the Slave Province, Canada.	230
Garanin, V.K., Kudriavtseva, G.P., Possukhova, T.V.	Diamonds of Arkhangelsk kimberlite province (review).	233
Garanin, V.K., Kudriavtseva, G.P., Vasilyeva, E.R.	The fundamental study of garnets: application for prospecting and economical estimation of diamond bearing kimberlites.	236
Gaspar, J.C., Araujo, D.P., Melo, M.V.L.C.	Olivine in Carbonatitic and Silicate Rocks in Carbonatite Complexes	239
Gaspar, J.C., Teixeira, N.A., Steele, I.M.	Cathodoluminescence of Juina Diamonds.	242
Gaul, O., O'Reilly, S.Y., Griffin, W.L.	Lithosphere Mapping in Eastern Australia.	245
Gelger, C.A.	Could the Effect of Order-Disorder in Garnet be Important for Upper Mantle Petrology?	248
Gibson, S.A., Thompson, R.N., Dickin, A.P.	Subcontinental mantle plume impact and kimberlite genesis.	250
Girnis, A.V., Stachel, T., Brey, G.P., Harris, J.W., Phillips, D.	Internally consistent geothermobarometers for garnet harzburgites.	253
Gonzaga, G.M., Gaspar, J.C., Araujo, D.P.	³ He and ¹⁰ Be Isotopes as a Diamond Exploration Tool: Some thoughts Based on Literature Data	256
Graham, I., Burgess, J.L., Bryan, D., Ravenscroft, P.J., Thomas, E., Doyle, B.J., Hopkins, R., Armstrong, K.A.	The Diavik Kimberlites - Lac de Gras, Northwest Territories, Canada.	259
Graham, S., Lambert, D.D., Shee, S.R., Smith, C.B., Hamilton, R.	Re-Os and Sm-Nd Isotope Systematics of Alkaline Ultramafic Rocks, Xenoliths and Macrocrysts from the Earaeedy Basin, Yilgarn Craton.	262
Graham, S., Lambert, D.D., Smith, Chris.B., Shee, S.R., Reeves, S.J.	Re-Os Isotope Systematics of Oxide Xenocrysts and Peridotite Xenoliths from the Kimberlites and the Argyle Lamproite, Kimberley Block, Australia: Implications for the Evolution of the Continental Lithospheric Mantle.	265
Greenwood, J.C., Gibson, S.A., Thompson, R.N., Weska, R.K., Dickin, A.P.	Petrogenesis of Cretaceous Kimberlites from the Paranatinga Region, Central Brazil.	268
Griffin, W.L., Doyle, B.J., Ryan, C.G., Pearson, N.J., O'Reilly, S.Y., Natapov, L., Kivi, K., Kretschmar, U., Ward, J.	Lithosphere Structure and Mantle Terranes: Slave Craton, Canada.	271
Griffin, W.L., Wln, T.T., Davies, R., Wathanakul, P., Andrew, A., Metcalfe, I.	Diamonds from Myanmar and Thailand: Characteristics and Possible Origins.	274
Grütter, H.S.	Chrome-calcium, Magnesium-number and Yttrium characteristics of garnets in depleted Lherzollitic, Harzburgitic and Dunitic mantles.	277
Grütter, H.S., Apter, D.B.	Kimberlite-and lamproite-borne chromite phenocrysts with "diamond-inclusion"-type chemistries.	280
Grütter, H.S., Apter, D.B.	Garnet xenocryst chemistries in a traverse from Eendekull to Kimberley over the south-western margin of the Kaapvaal craton.	283
Grütter, H.S., Quadling, K.E.	Some comments on the (ab)use of sodium in garnet to predict eclogitic diamond potential.	287
Gurney J.J., Moore R.O., Beil D.R.	Mineral associations and compositional evolution of Monastery kimberlite megacrysts	290
Haggerty, S.E., Fung, A.T.	Orbicular Oxides in Carbonatitic Kimberlites: High Pressure Autoliths or Low P Liquid Immiscibility?	293
Hamilton, M.A., Pearson, D.G., Stern, R.A., Boyd, F.R.	Constraints on MARID petrogenesis: SHRIMP II U-Pb zircon evidence for pre-eruption metasomatism at Kampfersdam.	296
Harlow, G.E.	Interpretation of Kcpx and CaEs in Clinopyroxene from Diamond Inclusions and Mantle Samples.	299
Harmer, R.E.	Carbonatite magmas in the mantle: Evidence and relationship to kimberlites, orangeites and lamproites.	302
Harris, P.D., Courtnage, P.M.	The effects of Regolith-Landform development in diamond exploration: spectral investigation.	305
Harte, B., Hutchison, M.T., Lee, M., Harris, J.W.	Inclusions of (Mg,Fe)O in Mantle Diamonds	308
Hatton, C.J.	The Difference between Sheared and Granular Peridotites.	311
Hatton, C.J.	The Kimberlite-Megacryst link at Monastery Mine.	314
Hauri, E.H., Pearson, D.G., Bulanova, G.P., Millledge, H.J.	Microscale variations in C and N isotopes within mantle diamonds revealed by SIMS.	317
Hausel, W.D., Kucera, R.E., McCandless, T.E., Gregory, R.W.	Mantle-Derived Diatremes in the Southern Green River Basin, Wyoming, USA	320
Heaman, L., Teixeira, N.A., Gobbo, L., Gaspar, J.C.	U-Pb Mantle Zircon Ages for Kimberlites from the Juina Paranatinga Provinces, Brazil.	322

Author	Title	Page No's
Hearman, L.M., Creaser, R.A., Cookenboo, H.O.	Zircons from eclogite in the Jericho Kimberlite Pipe, northern Canada: Evidence for Proterozoic High Pressure Metamorphism Beneath the Slave Province.	325
Hearn, B.C. Jr	Peridotite xenoliths from Porcupine Dome, Montana, USA: Depleted subcontinental lithosphere samples in an olivine-phlogopite-carbonate magma.	328
Helmstaedt, H.H., Harrap, R.M.	Tectonic Aspects of the Kimberlite - Diamond - Upper-Mantle-Sample Connection: Does a coherent Model evolve?	331
Hutchison, M.T., Cartigny, P., Harris, J.W.	Carbon and Nitrogen Compositions and Cathodoluminescence Characteristics of Transition Zone and Lower Mantle Diamonds from Sao Luiz, Brazil.	336
Ionov, D.A., Griffin, W.L., O'Reilly, S.Y.	Garnet peridotite xenoliths in alkali basalts from Siberia and Mongolia: a comparison of lithospheric mantle compositions in cratonic and younger terrains.	339
Iouchko, N.A., Kremenetsky, A.A., Kouznetsov, I.I.	Nature of Diamonds, Melts and Fluids in the Ring Structures: Endogeneous Explosion vs. Impact Process.	342
Irvine, G.J., Pearson, D.G., Carlson, R.W., Boyd, F.R.	Platinum Group Element Constraints On The Origin Of Cratonic peridotites: A Study Of Kimberley Peridotite Xenoliths.	346
Irving, A.J., Kuehner, S.M.	Petrology and Geochemistry of the Ruby Slipper Lamproite, Western Montana: A Leucite-Bearing, Ultrapotassic Magma in an Eocene Continental Arc.	349
Izraeli, E., Schrauder, M., Navon, O.	On the Connection between Fluid and Mineral-Inclusions in Diamonds.	352
Izraeli, E., Wilcock, I.C., Navon, O.	Raman Shifts of Diamond Inclusions - A Possible Barometer.	355
Jacob, D.E., Foley, S.F.	Evidence for Archean Ocean Crust with Island Arc Signature from Diamondiferous Eclogite Xenoliths.	358
Jacob, D.E., Kjarsgaard, B., Horn, I.	Trace element concentrations by Laser Ablation ICP-MS in subcalcic garnets from Saskatchewan and Somerset Island, Canada.	361
Jacob, D.E., Matthey, D.P.	Geochemistry of layered kyanite-bearing eclogites from the Roberts Victor Mine, South Africa.	364
James, D.E., van der Lee, S., Gao, S., Silver, P., VanDecar, J., Kuehnel, R., Jordan, T.H., Saitzer, R., Caherty, J., Gore, J., Zengeni, T., Nguun, T., Wright, C., Webb, S., Burford, D., Doucoure, M., Mollisana, M., Green, R., Robey, J., Harvey, J., Kostlin, E., Reichardt, F.	Review of Seismic Structure of the Continental Lithosphere with Results from the Southern Africa Seismic Experiment.	366
Janney, P.E., le Roex, A.P.	Causes of Compositional Diversity in the Olivine Melilitites of Namaqualand-Bushmanland, South Africa.	371
Janney, P.E., le Roex, A.P., Viljoen, K.S.	Trace Element and Isotopic Characteristics of Olivine Melilitites from the Western Cape, South Africa: Implications for the Sources of Group 1 Kimberlites.	374
Janse, A.J.A.	Archons, Protons and Tectons: an update.	377
Johnson, L.H., Burgess, R., Turner, G., Milledge, H.J.	Fluids trapped within diamond: clues to mantle geochemistry.	380
Johnson, L.H., Burgess, R., Turner, G., Milledge, H.J.	Noble gas and halogen systematics of fluids within diamond coats from Canada and Africa.	383
Jones, A.P., Dobson, D., Milledge, H.J., Taniguchi, T., Litvin, Y., Genge, M.J., Rabe, R.	Experiments with low-T potassic carbonatitic melts, fluids and diamonds.	386
Kaminsky, F.V., Gorzynsky, G., Sablukova, L.I., Sablukov, S.M., Zakharchenko, O.D.	Primary Sources of Diamonds in the Birim Area, Ghana.	389
Kaminsky, F.V., Sablukov, S.M., Sablukova, L.I., Shpanov, V.E., Zhuravlev, D.Z.	Diamondiferous Minette Dykes from the Parker Lake Area, N.W.T., Canada.	392
Kaminsky, F.V., Zakharchenko, O.D., Channer, D.M., DeR., Blinova, G.K., Maltsev, K.A.	Diamonds from the Guaniamo area, Venezuela.	395
Kelemen, P.B.	One view on the genesis of cratonic mantle peridotites.	398
Keller, R.A., Remley, D.A., Snyder, G.A., Taylor, L.A., Sobolev, N.V.	Mantle Xenoliths from the Obnazhennaya Kimberlite, Yakutia.	402
Keller, R.A., Taylor, L.A., Snyder, G.A., Sobolev, V.S., Carlson, W.D., Sobolev, N.V., Pokhilenko, N.P.	3-D Petrography of a Diamondiferous Eclogite from Udachnaya, Siberia.	405
Kempton, P.D., Hawkesworth, C.J., Lopez-Escobar, L., Ware, A.J.	Geochemistry of spinel ± garnet lherzolite xenoliths from Pali Aike: Implications for evolution of mantle lithosphere beneath southern Patagonia	408
Kent, R.W., Paul, D.K., Basu, A., Ghose, N.C., Kempton, P.D.	Mafic alkaline intrusions in the Damodar Valley, India: the micaceous kimberlite-lamproite connection revisited.	411
Kepezhinskas, K., Kepezhinskas, P.	Ultramafic - Mafic rocks of the Eastern European Craton and Their Diamond Potential.	414
Kepezhinskas, P., Defant, M.J., Maury, R., Clague, A., Joron, J.-L., Cotten, J.	Composition of Island-Arc Mantle and its Bearing on the Origin of Cratonic Lithosphere.	417
Kilty, S.J.	Airborne Electromagnetic and Magnetic Surveying in the Search for Kimberlites.	420
Kinny, P.D., Trautman, R.L., Griffin, B.J., Harte, B.	Carbon isotopic analyses of microdiamonds.	423
Kinzler, R.J., Grove, T.L.	Origin of Depleted Cratonic Harzburgite by Deep Fractional Melt Extraction and Shallow Olivine Cumulate Infusion.	426

Author	Title	Page No's
Kirkley, M.B., Kolebaba, M.R., Carlson, J.A., Gonzales, A.M., Dyck, D.R., Dierker, C.	Kimberlite Emplacement Processes Interpreted from Lac de Gras Examples.	429
Kivilets, G., Phillips, D., Shee, S.R., Vercoe, S.C., Barton, E.S., Smith, C.B., Fourie, L.F.	$^{40}\text{Ar}/^{39}\text{Ar}$ Dating of yimengite from the Turkey Well kimberlite, Australia: The oldest and the rarest.	432
Kjarsgaard, B.	Compositional trends of spinel and mica in alkali minettes, southern Alberta, Canada.	435
Klemme, S., O'Neill, H.St.C.	The partitioning of Chromium between orthopyroxene and spinel in the system $\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{Cr}_2\text{O}_3$: Implications for geothermobarometry for Upper Mantle rocks.	438
Klump, J., Gurney, J.J.	A Pilot Study of the Swartruggens Kimberlite Dyke Swarm.	441
Koga, K.T., Shimizu, N., Grove, T.L.	Dis-equilibrium trace element re-distribution during garnet to spinel facies transformation.	443
Kong, J.M., Boucher, D.R., Scott-Smith, B.H.	Exploration and Geology of the Attawapiskat kimberlites, James Bay Lowland, Northern Ontario, Canada.	446
Kopylova, M.G., Russell, J.K., Cookenboo, H.	Petrography and Chemistry of the Jericho kimberlite (Slave Craton, Northern Canada).	449
Kopylova, M.G., Russell, J.K., Cookenboo, H.	Upper mantle stratigraphy and thermal regime of the north central Slave Craton, Canada.	452
Kopylova, M.G., Russell, J.K., Cookenboo, H.	Unique chemical features of the peridotitic mantle below the Jericho kimberlite (Slave Craton, Northern Canada).	455
Kornilova, V.P., Safronov, A.F., Philippov, N.D., Zayzev, A.I.	The garnet of diamond association in lamprophyres from the Anabar massif.	458
Kostrovitsky, S.I., Pavlova, L.A., Suvorova, L.V.	Preliminary information about the first finding of Ti-bearing kirschsteinite (Ferromonticellite) in kimberlites.	460
Kostrovitsky, S.I., de Bruin, D.	Ultramafic association of minerals (garnet-ureyite diopside-chromspinel) in micaeous kimberlites of Yakutian province.	463
Kostrovitsky, S.I., Morikyo, T.	Sr,Nd isotopic data of kimberlites and related rocks from North of Yakutian kimberlite province (Russia).	466
Kouznetsova, E.I., Galdin, N.E.	Continental lithosphere deep structure researches on the base of scientific deep drilling	469
Kravchenko, S.M.	Kimberlite Types IA, IB, and II as Series from Different Mantle Depths.	471
Kryvoslyk, I.N.	Brief Review of the Theory of Liquid Immiscibility of Kimberlite Magma.	473
Kuehner, S.M., Irving, A.J.	Corundum-kyanite Eclogite, Grosspyrite and Epidote Amphibolite of Probable Subducted Slab Origin from Paleogene Diamondiferous Pipes in SW Wyoming	475
Kukkonen, I.T., Peitonen, P.	Geotherm and a rheological profile for the central Fennoscandian lithosphere.	478
Kuligin, S.S., Pokhilenko, N.P.	Mineralogy of xenoliths of garnet pyroxenites from kimberlite pipes of Siberian Platform.	480
Kurszlauskis, S., Lorenz, V., Zimanowski, B., Büttner, R.	Experiments on explosive interaction of molten kimberlite with injected water.	483
LeCheminant, A.N., Heaman, L.M., Kretschmar, U., LeCouteur, P.C.	Complex Origins and Multiple Ages of Mantle Zircon Megacrysts from Canadian and South African Kimberlites.	486
Lee, C.T.	Are infected geotherms real?	489
Lee, C.T., Rudnick, R.L.	The origin and demise of cratonic lithosphere: a geochemical perspective from the Tanzanian craton.	492
Leggatt, P.B., Klinkert, P.S.	The application of Airborne Electromagnetic methods in the search for buried Kimberlites and Diamondiferous Gravels.	495
Leluyh, M.I., Kostrovitsky, S.I., Bezborodov, S.M., Nikulin, V.I., Prokopen, S.A., Serov, V.P., Tolstov, A.V., Zuev, V.M.	Kimberlites and related rocks of Anabar region (Yakutia, Russia).	497
Letendre, J., McCandless, T.E., Eastoe, C.J.	Morphology and Carbon Isotope Composition of Microdiamonds from Dachine, French Guiana.	500
Litasov, K.D., Kostrovitsky, S.I., Litasov, Yu.D.	Comparison of ilmenite-clinopyroxene symplectites from Vitim alkaline basalts and Yakutian kimberlites (Siberia, Russia).	503
Litasov, K.D., Litasov, Yu.D.	Reactional and differentiated pyroxenite xenoliths from alkaline basalts of the Vitim volcanic field (East Siberia): their role in metasomatism and position in mantle magmatic system.	506
Litasov, Yu.D., Nilda, K., Litasov, K.D.	Reactional modification of the primitive mantle by basaltic melts: an evidence from mantle-derived xenoliths of the Vitim Plateau (Russia).	509
Logvinova, A.M., Fedorova, E.N., Sobolev, N.V.	Microdiamonds from the Yubileynaya kimberlite pipe, Yakutia: morphology, physical properties, and mineral inclusions	512
Luk'yanova, L.I., Lobkova, L.P., Zhukov, V.V., Rybačchenko, A.Y., Ostroumov, V.P.	Diamonds of the Uralis Mobile Belt and Source Rocks for the Uralian (Brazilian) type Diamond Placers.	515
Lütjen, H., Blume, J., Pretorius, C.C.	Geophysical survey over the Elizabeth Bay Mine, Namibia.	518
Mabuza, M., Viljoen, K.S., Majola, S.	New diamond-bearing xenoliths from the Orapa Mine, Botswana.	521
Machin, K.J., Barton E.S.	The petrology of the Rex Mine kimberlite fissures, central Free State, South Africa.	524
Magee, C.W., Taylor, W.R.	Constraints on the history and origin of carbonado from luminescence studies.	527
Mahotkin, I.L.	Petrology of Group 2 Kimberlite - Olivine lamproite (K2L) from the Kostomuksha area, Karelia, N.W. Russia.	529
Mahotkin, I.L., Skinner, E.M.W.	Kimberlites from the Archangelsk region - A rock type transitional between kimberlites, melnoites and lamproites.	532
Mainprice, D., Barnuel, G., Ben Ismail, W., Lloyd, G.	Automatic crystal orientation mapping of Kimberlite nodules using electron back scattered diffraction in the scanning electron microscope.	535

Author	Title	Page No's
Mal'kov, B.A.	The Cosmic Cycles of Kimberlite Volkanism: New Data.	537
Mal'kov, B.A., Malyshev, N.A.	Diamond Occurrences in Kimberlites and Lamproites from Phanerozoic Mobile Belts on Example of the Timans, Urals and Ouachita.	540
Maikovets, V.G., Ionov, D.A., Griffin, W.L., O'Reilly, S.Y., Pokhilenko, N.P., Litasov, K.D.	A-P-T-composition cross-section of spinel and garnet facies lithospheric mantle in the Minusa region SW of the Siberian craton.	543
Marakushev, A.A., Bobrov, A.V.	Crystallization of Eclogite and Pyroxenite Magmas in the Diamond Depth Facies: Evidence from Garnet-Clinopyroxene Association.	546
Mason, P.R.D., Downes, H., Jarvis, K., Vannucci, R.	An Investigation of Incompatible trace elements in Massif Central mantle xenoliths by laser ablation ICP-MS: a new tool for investigating mantle geochemistry.	549
Massonne, H.J.	A new occurrence of microdiamonds in quartzofeldspathic rocks of the Saxonian Erzgebirge, Germany, and their metamorphic evolution.	552
McCammom, C.A.	Methods for Determination of Fe ³⁺ /SFe in Microscopic Samples.	555
McCandless, T.E.	Kimberlites: the Products of Deep-Seated Subduction.	558
Mc Dade, P., Harris, J.W.	Syngenetic Inclusion bearing diamonds from Letseng-la-Terai, Lesotho.	561
McKinlay, F.T., Scott Smith, B.H., de Gasparis, S., Kong, J.	Geology of the Recently Discovered Hardy Lake Kimberlites, NWT	564
Mendelsohn, M., Milledge, H.J.	Characterisation of diamonds by infrared spectroscopy.	567
Menzies, A.H., Baumgartner, M.C.	Application of garnet geothermobarometry to southern African kimberlites.	570
Menzies, A.H., Gurney, J.J., Harte, B., Hauri, E.	REE patterns in diamond bearing eclogites and diamond bearing peridotites from Newlands Kimberlite	573
Menzies, A.H., Milledge, H.J.M., Gurney, J.J.	Fourier Transform Infra-red (FTIR) Spectroscopy of Newlands diamonds	576
Menzies, A.H., Shirey, S.B., Carlson, R.W., Gurney, J.J.	Re-Os isotope systematics of diamond-bearing eclogites and peridotites from Newlands Kimberlite	579
Mikhailov, M.V., Kuznetsova, M.Yu., Kuzmina, T.S., Polyakov, A.A., Lukyanova, L.I.	New data on potential diamond presence in Western Russia.	582
Milashay, V.A.	Energy of kimberlite formation.	584
Milledge, H.J., Sutherland, F.L., Kennewell, P.	Further studies of Copeton Diamonds.	587
Milledge, H.J., Woods, P.A., Beard, A.D., Shelkov, D., Willis, B.	Cathodoluminescence of polished carbonado.	589
Miller, A.R., Sellar, M.H., Armitage, A.E., Davis, W.J., Barnett, R.L.	Late Triassic kimberlitic magmatism, western Churchill Structural Province, Canada.	591
Milstein, E.D., Erinchek, Yu.M., Egorkin, A.V., Parasotka, B.S.	The structure of the Lithosphere in Diamond-Bearing Kimberlite Areas. The Siberian Platform.	594
Mironov, V.P.	Internal Morphology of Diamonds from Pipe Udachnaya According to the Data of Luminescence Tomography Method.	597
Mitchell, R.H., Scott Smith, B.H., Larsen, L.M.	Mineralogy of Ultramafic Dikes from the Sarfartoq, Sisimiut and Manilsoq areas, west Greenland: Kimberlites or Melnoites?	600
Mitioukhine, S.I.	Chief Feature of Rocks of the Earth's crust within Kimberlite Provinces - Moderation of their Petrochemical Indicators.	603
Mitioukhine, S.I., Manakov, A.V., Pollaratskaya, O.L., Romanov, N.N.	New Data about the Structure of the Earth's crust according to Regional Geophysical Investigations: Results within Yakutian Kimberlitic Subprovince.	606
Moser, D.E., Hart, R.J.	Neoproterozoic and Paleoproterozoic re-activation of the crust-mantle transition beneath the central Kaapvaal craton, Laca kimberlite.	609
Nassichuk, W.W., Dyck, D.R.	Fossils Recovered from Kimberlite Pipes in the Lac de Gras Field, Slave Province, Northwest Canada; Geological Implications.	612
Natapov, L., Griffin, W.L.	Geodynamic controls on the distribution of diamondiferous kimberlites.	615
Navon, O.	Diamond formation in the Earth's mantle	618
Nguno Muatara, A.	Indicator minerals in kimberlites and their respective stream sediments, Gibeon Kimberlite Province, Namibia.	622
Nixon, P.H., Pearson, D.G.	Ultra-magnesian komatites of phanerozoic age, from SE Spain.	625
Nowell, G.M., Kempton, P.D., Pearson, D.G.	Hf-Nd Isotope Systematics of Kimberlites: Relevance to Terrestrial Hf-Nd systematics.	628
Nowell, G.M., Kempton, P.D., Pearson, D.G.	Trace Element and Isotope Geochemistry of Siberian Kimberlites.	631
Nowell, G.M., Pearson, D.G.	Hf Isotope Constraints on the Genesis of Kimberlitic Megacrysts: Evidence for a Deep Mantle Component in Kimberlites.	634
Nowell, G.M., Pearson, D.G., Kempton, P.D., Irving, A.J., Turner, S.	A Hf Isotope Study of Lamproites: Implications for their Origins and Relationship to Kimberlites	637
Nowell, G.M., Pearson, D.G., Kempton, P.D., Noble, S.R., Smith, C.B.	The source regions/components of kimberlites: Constraints from Hf-Nd isotope systematics.	640
O'Brien, H.E., Tyni, M.	Mineralogy and geochemistry of Kimberlites and related rocks from Finland.	643
O'Reilly, S.Y., Griffin, W.L., Poudjorn Djomani, Y.	Are Lithospheres forever?	646
Pal'yanov, Yu.N., Gusev, V.A., Kupriyanov, I.N., Borzdov, Yu.M., Sokol, A.G., Khokhriakov, A.F., Sobolev, N.V.	The effect of growth rate on formation of nitrogenous defects in diamond.	649
Panina, L.I.	Genesis of Cocites from North Vietnam: Results of Melt Inclusions Studies in Minerals.	652
Panina, L.I., Usoltseva, L.M.	The Role of High-Calcium Alkaline Sulfate-Carbonate Melts in Formation of Melilitite-Monticellite Rocks and Carbonatites of the Malyi Murun Massif (Aldan, Russia).	655

Author	Title	Page No's
Pearson, D., Carlson, R.W., Boyd, F.R., Shirey, S.B., Nixon, P.H.	Lithospheric mantle growth around cratons: A Re-Os isotope study of peridotite xenoliths from East Griqualand.	658
Pearson, D.G., Shirey, S.B., Bulanova, G.P., Carlson, R.W., Milledge, H.J.	Dating diamonds using the Re - Os isotope technique: A study of sulfide inclusions in Siberian diamonds.	661
Pearson, D.G., Davies, R.M., Shirey, S. B., Carlson, R.W., Griffin, W.L.	The age and origin of eastern Australian diamonds: Re-Os isotope evidence from sulfide inclusions in two diamonds from Wellington, New South Wales.	664
Pearson, D.G., Milledge, H.J.	Diamond growth conditions and preservation; Inferences from trace elements in a large garnet inclusion in a Siberian diamond.	667
Pearson, N.J., Griffin, W.L., Doyle, B.J., O'Reilly, S.Y., van Achterbergh, E., Kivi, K.	Xenoliths from kimberlite pipes of the Lac de Gras area, Slave Craton, Canada.	670
Pearson, N.J., Griffin, W.L., Kaminsky F.Y., van Achterbergh, E., O'Reilly, S.Y.	Trace element discrimination of garnet from diamondiferous kimberlites and lamproites.	673
Peltonen, P.	Silicification of Garnet Peridotite Xenoliths from the Lahtojoki Kimberlite Pipe, Eastern Finland.	676
Peltonen, P., Huhma, H., Tyni, M., Shimizu, N.	Garnet Peridotite Xenoliths from Kimberlites of Finland: Nature of the Lithospheric Mantle at Archaean craton - Palaeoproterozoic mobile belt transition.	678
Pendock, N.	Breaking the Nyquist Barrier: Superresolution Magnetic Imaging from Gradient Data.	681
Perelra, R.S., Wheelock, G., Bizzi, L., Silva, H., Leite, A.	Alluvial diamond potential of Paleo-Drainage systems in the Headwaters of the Sao Francisco River, Minas Gerais, Brazil	684
Phillips, D., Harris, J.W., Kiviets, G.B., Burgess, R., Fourie, L.F.	40Ar/39Ar Laser Probe Analyses of Clinopyroxene Diamond inclusions from the Orapa and Mbuyi-Miya Mines.	687
Phillips, D., Kiviets, G.B., Barton, E.S., Smith, C.B., Viljoen, K.S., Fourie, L.F.	40Ar/39Ar Dating of Kimberlites and Related Rocks: Problems and Solutions.	690
Pizzolato, L.A., Schulza, D.J.	Preliminary investigations of megacrysts and peridotite xenoliths from the Kelsey Lake kimberlite, Colorado-Wyoming, USA.	693
Podvysotsky, V.T., Zuev, V.M., Nikulin, V.I., Lelyoukh, M.I., Bezborodov, S.M.	Conception of Formation of Magmatogene and Terrigenous Diamondiferous Formations of Ancient Platforms as the basis of deposits' forecast.	696
Pokhilenko, N.P., McDonald, J.A., Metnyk, W., Hall, A.E., Shimizu, N., Vavilov, M.A., Afanasiev V.P., Reimers, L.F., Irvin, J., Pokhilenko, L.N., Vasilenko, V.B., Kulgin, S.S., Sobolev, N.V.	Kimberlites of Camsell Lake field and some features of construction, and composition of lithosphere: roots of southeastern part of Slave Craton, Canada.	699
Pokhilenko, N.P., Sobolev, N.V., Kulgin, S.S., Shimizu, N.	Peculiarities of pyroxenite paragenesis garnets distribution in Yakutian kimberlites and some aspects of the Siberian craton lithospheric mantle evolution.	702
Presnall, D.C., Walter, M.J.	High pressure phase equilibrium constraints on the origin of eclogites.	705
Pretorius, C.C., Blume, J., Lütjen, H., Trofimczyk, K.	Results of Geophysical trials to profile the Kimberlite/Host rock contacts at Venetia Mine and the BK-9 Pipe.	708
Pretorius, W., Barton, J.M., Jr.	The Use of Amphibolite Melting Experiments in Constraining Conditions of Melting in Natural Amphibolite Nodules from the Venetia Kimberlite Pipes.	710
Pretorius, W., Leahy, K.	Implications for diamond prospectivity from comparisons of diamond-bearing lithosphere in two Proterozoic orogenic belts.	713
Prikhodko, V.S., Zemlyanukhin, V.N.	Petrology of spinel Peridotite Xenoliths from Cenozoic Basaltoids in the Khanka Craton's Terrain (East Russia)	716
Pybus, G.Q.J., Hussey, M.C., Linton, P.L.	Spectral investigations of a variety of magnesium-bearing rock types: Implications for kimberlite exploration.	717
Rapp, R.P., Shimizu, N.	Subduction and Slab Melting in the Archean: Experimental Constraints and Implications for Development of Cratonic Lithosphere.	720
Rass, I.T., Gerasimov, V.Yu., Laputina, I.P., Ilupin, I.P.	Diamond occurrence in kimberlites dependent on melting depths and rates of cooling of parental mantle magmas.	723
Rass, I.T., Kravchenko, S.M.	Melilite-Bearing Rocks within Alkaline-Ultrabasic Complexes: Derivatives from SiO ₂ -Poor, Ca-Rich Mantle Magma?	725
Reid D.L., Cooper A. F.	Carbonatite and silicate magmas at Dicker Willem, southern Namibia : their origin and source region characteristics	727
Reimers, L.F., Pokhilenko, N.P., Yefimova, E.S., Sobolev, N.V.	Ultramafic mantle assemblages from Sytykanskaya kimberlite pipe (Yakutia).	730
Rice, A.	Can the blasting excavation engineering sciences provide insight into the processes of kimberlite emplacement and eruption?	733
Richardson, S.H., Chinn, I.L., Harris, J.W.	Age and Origin of Eclogitic Diamonds from the Jwaneng Kimberlite, Botswana.	734
Robinson, D.N., Ferraris, R., Anderson, V.G., Parker, G.M., Van Blerck, E., Hart, D.	Colour, Morphological and Surface Textural Characteristics of the Diamonds in Venetia Kimberlites.	737
Roden, M.F., Laz'ko, E.E., Jagoutz, E.	Petrology and Geochemistry of Peridotite Inclusions from the Mir kimberlite, Siberia	740
Rodionov, A.S., Viljoen, K.S.	Venetia megacrysts, Northern Province, South Africa.	743
Romashkin, A.I.	Potassium Alkaline Magmatism in the Russian Far East.	746
Romashkin, A.I., Kukhtina, L.M.	Mineralogy of Inggilite.	749
Rombouts, L.	Extreme Value Analysis of Diamond Sizes and Values.	752
Rudnick, R.L., Ireland, T.R., Gehrels, G., Irving, A.J., Chesley, J.T., Hanchar, J.M.	Dating Mantle Metasomatism: U-Pb Geochronology of zircons in cratonic mantle xenoliths from Montana and Tanzania.	754
Ruiz, J., McCandless, T.E., Helmstaedt, H.H.	Eclogites from the Colorado Plateau: A Phanerozoic Record of Subduction beneath North America.	757

Author	Title	Page No's
Saltykov, O.G., Erinchek, Yu.M., Milshtein, E.D	The model of buried kimberlite field based on its reflection in postkimberlite reservoir rocks (by the example of the Yakutian province).	760
Saparin, G.V., Obyden, S.K., Titkov, S.V	Use of Cathodoluminescence Scanning Electron Microscope with Color TV Display for Study of Natural Diamonds Internal Structure.	763
Schmitz, M.D., Bowring, S.A., Robey, J.V.A.	Constraining the thermal history of an Archean craton: U-Pb thermochronology of lower crustal xenoliths from the Kaapvaal craton, southern Africa	766
Schulze, D.J., Valley, J.W., Bell, D.R., Spicuzza, M.	Significance of Oxygen Isotope Variations in the Cr-Poor Megacryst Suite.	769
Scott Smith, B.H., Orr, R.G., Robertshaw, P., Avery, R.W.	Geology of the Fort A La Corne Kimberlites, Saskatchewan.	772
Seggie, A.G., Hannweg, G.W., Colgan, E.A., Smith, C.B.	The Geology and Geochemistry of the Venetia Kimberlite cluster, Northern Province, South Africa.	775
Seitz, H.M., Woodland, A.B	Lithium and Beryllium abundances in peridotitic, pyroxenitic and eclogitic mantle assemblages.	778
Seife, G.R., Trofimczyk, K.K.	Recent developments in the application of Borehole Geophysical logging techniques in Diamond Mining and exploration - some case studies.	781
Shamshina, E.A., Zaitsev, A.I.	New age of Yakutian kimberlites.	783
Sharygin, V.V	Melt inclusions and Chromite In Lamprolites from Smoky Butte, Montana.	785
Sharygin, V.V., Golovin, A.V., Smirnov, S.Z., Malkovets, V.G	Relationship between Websterite Xenolith and Host Basanite (Pipe Bele, Khakasia, Russia): Evidence from Fluid and Silicate-Melt Inclusions in Minerals.	788
Sharygin, V.V., Litasov, K.D., Smirnov, S.Z., Kuzmin, D.V., Reutsky, V.N., Ivanov, A.V	Fluid and Silicate-Melt Inclusions and Interstitial Glass in Mantle Xenoliths from Melanephelinites of the Udokan Lava Plateau, Russia.	791
Sharygin, V.V., Pospelova, L.N	Sulphide inclusions in Early Lamprolite Minerals.	794
Shatsky, V.S., Zedgenizov, D.A., Yefimova, E.S., Rylov, G.M., De Corte, K., Sobolev, N.V.	A comparison of morphology and physical properties of microdiamonds from the mantle and crustal environments.	797
Shee, S.R., Vercoe, S.C., Wyatt, B.A., Campbell, A.N., Colgan, E.A., Hwang, P.H.	Discovery and Geology of the Nabberu Kimberlite Province, Western Australia.	800
Shimizu, N., Pokhilenko, N.P., Boyd, F.R., Pearson, D.G	Trace element characteristics of garnet dunites/harzburgites, host rocks for Siberian peridotitic diamonds.	803
Shimizu, N., Pokhilenko, N.P., McDonald, J.A	Geochemical characteristics of the Slave craton lithosphere: A view from heavy mineral concentrate garnets.	805
Shimizu, N., Sobolev, N.V., Yefimova, E.S.	Trace element heterogeneities in-situ diamond inclusion garnets from Siberia	807
Shirey, S.B., Carlson, R.W., Gurney, J.J., van Heerden, L.	Re-Os Isotope Systematics of Eclogites from Roberts Victor: Implications for Diamond Growth and Archean Tectonic Processes.	808
Shiryaev, A.A., Galimov, E.M., Sobolev, N.V., Kolesov, G.M., Zakharchenko, O.D.	Trace Elements in Inclusion-free Diamonds from Venezuela and Arkhangelsk Deposits.	811
Simakov, S.K.	Garnet - clinopyroxene geobarometry of deep mantle eclogites and eclogite paleogeotherm	814
Skinner, E.M.W., Mahotkin, I.L., Grütter, H.S	"Mellite" in Kimberlites.	817
Smith, S.C., Ihinger, P.D.	Geochemical evolution of the New England lamprophyre suite: a hotspot signature preserved in the continental crust?	820
Snyder, G.A., Keller, R.A., Taylor, L.A., Remley, D., Halliday, A.N., Sobolev, N	The Origin of Ultramafic (Group A) Eclogites: Nd & Sr Isotopic Evidence From the Obnazhennaya Kimberlite, Yakutia.	823
Snyder, G.A., Taylor, L.A., Beard, B.L., Halliday, A.N., Sobolev, N.V	The Diamond-Bearing Mir Eclogites, Yakutia: Nd And Sr Isotopic Evidence for Continental Crustal Input In An Archean Oceanic Environment.	826
Sobolev, N.V., Yefimova, E.S., Channer, D., Anderson, P.F.N., Barron, K.M	A unique eclogitic source of Guaniamo diamonds, Guyana shield, Venezuela.	829
Sobolev, N.V., Yefimova, E.S., Koptil, V.I	Crystalline Inclusions in Diamonds in the Northeast of the Yakutian Diamondiferous Province.	832
Sobolev, V.N., Taylor, L.A., Snyder, G.A., Jerde, E.A., Neal, C.A., Sobolev, N.V	Metasomatism of the Mantle Beneath Yakutia: A Quantitative Study of Secondary Chemistry and Mineralogy in Udachnaya Eclogites.	835
Sokolovsky, A.K., Serokurov, Yu.N., Kalmykov, V.D	System analysis of remote sensing data on structural control of diamondiferous areas	838
Solovjeva, L.V., Barankevich, V.G., Bayukov, O.A., Glazunov, O.M	Polychrome olivines in coarse grained lherzolites from the Udachnaya pipe - possible fine indicators of reduced metasomatism	841
Spetsius, Z.V	Two Generations of Diamonds in the Eclogite Xenoliths.	844
Spetsius, Z.V., Bezborodov, S.M	Compositional variations and floatability of Kimberlite Ores of Russia	847
Spetsius, Z.V., Griffin, B.J	Secondary phases associated around diamonds in eclogites from the Udachnaya kimberlite pipe: Implications for diamond genesis.	850
Spetsius, Z.V., Griffin, W.L	Trace element composition of garnet kelyphites in xenoliths from Udachnaya as evidence of their origin.	853
Spetsius, Z.V., Taylor, W.R., Griffin, B.J.	Major and trace element partitioning between mineral phases in diamondiferous and non-diamondiferous eclogites from the Udachnaya kimberlite pipe, Yakutia.	856
Stachel, T., Harris, J.W., Brey, G.P	Inclusions in diamonds from Mwadui (Tanzania) - chemical mush in the source.	859
Stachel, T., Viljoen, K.S., Harris, J.W., Brey, G.P	REE patterns of garnets from diamonds and diamondiferous peridotites - geochemical signatures of the diamond source.	862
Stasiuk, L.D., Lockhart, G.D., Nassichuck, W.W., Carlson, J.A., Tomica, M.	Kimberlite emplacement temperatures derived from the thermal history of Organic matter, Lac de Gras, Canada.	865

Author	Title	Page No's
Stiefenhofer, J., Viljoen, K.S., Tainton, K.M., Dobbe, R., Hannweg, G.W.	The petrology of a mantle xenolith suite from Venetia, South Africa.	888
St Pierre, M., Wynne, P.J., Counts, B.	Paleomagnetisation of Kimberlites on the BHP/Dia Met Diamond Project.	874
Sweeney, R.J., Konzett, J., Prozesky, V.M.	The determination of hydrogen in peridotitic minerals by nuclear methods.	874
Sweeney, R.J., Winter, F.	Kimberlite as high-pressure melts: the determination of segregation depth from major element chemistry.	877
Tainton, K.M., Seggie, A., Bayly, B.A., Tomlinson, I., Quadling, K.E.	Regional variation in mantle heat flow within the Tanzanian Craton.	880
Taylor, L.A., Bulanova, G.P., Snyder, G.A., Keller, R.A.	Multiple Inclusions In Diamonds: Evidence For Complex Petrogenesis.	883
Taylor, W.R., Bristow, J.	Cyclicality of Continental Alkaline Magmatism in the Geological Record.	886
Taylor, W.R., Jaques, A.L.	Crystallization history of the Argyle and Ellendale olivine lamproites: constraints from spinel-olivine thermometry and oxygen barometry.	888
Taylor, W.R., Kamperman, M., Hamilton, R.	New thermometer and oxygen fugacity sensor calibrations for ilmenite- and chromian spinel-bearing peridotitic assemblages.	891
Taylor, W.R., Kingdom, L.	Mineralogy of the Jagersfontein kimberlite - an unusual Group I micaceous kimberlite - and a comment on the robustness of the mineralogical definition of 'orangeite'.	892
Taylor, W.R., Matveev, S.	Recalibration of the 5-parameter MRK equation of state for C-O-H fluids under upper mantle conditions and some experimental tests.	895
Taylor, W.R., Nimis, P.	A single-pyroxene thermobarometer for Iherzolitic Cr-diopside and its application in diamond exploration.	897
Taylor, W.R., Reddcliffe, T.H., Jakimowicz, J.	Thermobarometry of peridotitic Cr-diopside from the Merlin kimberlites, Northern Territory, Australia - nature of the upper mantle beneath the Proterozoic North Australian craton.	899
Teixeira, N.A., Gaspar, J.C., Oliveira, A.L.A.M., Bitencourt, R.M., Yeda, B.	Morphology of the Juina Maars	902
Teixeira, N.A., Gaspar, J.C., Waissel, O., Almeida, A.J., Bellther, J.A., Gobbo, L.	Geology of the Juina Diamondiferous Province	905
Thomas, R.D., Novak, N.A., Janse, A.J.A.	Diamonds in ultrabasic rock near Wawa, Ontario, Canada.	908
Titkov, S.V., Bershov, L.V., Scandale, E., Saparin, G.V., Chukichev, M.V., Speranskiy, A.V.	Nickel Structural Impurities in Natural Diamonds.	911
Titkov, S.V., Gorshkov, A.I., Vinokurov, S.F., Bershov, L.V., Solodov, D.I., Sivtsov, A.V.	Carbonado from Yakutian diamond deposits (Russia): microinclusions, impurities, and paramagnetic centers.	914
Tompkins, L.A., Meyer, S.P., Han, Z., Hu, S.	Petrology and Geochemistry of Kimberlites from Liaoning and Shandong Provinces, China.	917
Tompkins, L.A., Taylor, W.R., Ramsay, R.R., Armstrong, R.	The Mineralogy and Geochemistry of the Kamafugitic Tres Barras Intrusion, Mata da Corda, Minas Gerais, Brazil.	920
Tomshin, M.D., Fomin, A.S., Oleinikov, B.V.	Basites of the Vilyui - Markha zone (Siberian platform).	923
Trautman, R.L., Griffin, B.J., Bulanova, G.P.	Growth features and nitrogen aggregation properties of microdiamonds derived from kimberlite diatremes.	926
Tsyganov, V.A., Kontarovich, R.S.	Target-specific airborne geophysical forecast-exploration technology for diamond deposits <Field - Cluster - Pipe>.	929
Ulmer, G.C., Grandstaff, D.E., Göbbels, M., Woermann, E.	An Experimental Delineation of the Oxygen Fugacity of Moissanite (SiC) bearing Silicate Systems	932
Van Acherterbergh, E., Griffin, W.L., Shee, S.R., Wyatt, B.A., Sharma, A.L.	Natural Trace Element Distribution Coefficients for Garnet, Clino- and Orthopyroxene: Variations with Temperature and Pressure.	934
Van Acherterbergh, E., Griffin, W., Stiefenhofer, J.	Xenoliths from the Letlhakane Kimberlite: Geochemistry and Implications for Mantle Processes.	937
Verichev, E.M., Sablukov, S.M., Sablukova, L.I., Zhuravlev, D.Z.	A new type of diamondiferous kimberlite of the Zimny Bereg area (pipe named after Vladimir Grib).	940
Viljoen, K.S., Phillips, D., Harris, J.W., Robinson, D.N.	Mineral inclusions in diamonds from the Venetia kimberlites, Northern Province, South Africa.	943
Vladykin, N.V., Lelyukh, M.I., Tolstov, A.V.	Lamproites of the Anabar region, Northern rimming of the Siberian platform.	946
Voinova, I.P., Prikhodko, V.S.	Post-accretionary stage in the evolution of ultramafic magmatism in accretionary prisms: rock types, diamond potential (on example of East Russia).	949
Vouiko, V.	Method of Quantitative Evaluation of Kimberlite Pipes' Productivity.	950
Ward, J.R., Norman, D.I.	Geochemical and Physical Aspects of Diamonds from the Akwatia and Tarkwa Diamondfields in Southern Ghana, West Africa	953
Williams, C.M., Robey, J.v.A., Abson, J.P.	Petrography and Mineral Chemistry of the Mwenezi-01 Kimberlites, Zimbabwe.	955
Woermann, E., Göbbels, M., Ulmer, G.C., Grandstaff, D.E.	Moissanite and its bearing on the oxygen fugacity of the deeper regimes of the Earth's upper mantle	958
Wood, B.D., Scott Smith, B.H., de Gasparis, S.	The Mountain Lake Kimberlitic Pipes of NW Alberta: Exploration, Geology and Emplacement Model.	960
Woodland, A.B., Peltonen, P.	Ferric/Ferrous Iron Contents of Garnet and Clinopyroxene and Calculated Oxygen Fugacities of Peridotite Xenoliths from the Eastern Finland Kimberlite Province.	963
Wyatt, B.A., Ma Wenyun, Li Ziyun, Joyce, J., Colgan, E.A., Smit, D., De Bels, M.	The Ningxiang Lamproites, Hunan Province, China: Petrology and Mineral Chemistry.	965
Wyatt, B.A., Morfi, L., Gurney, J.J., Pearson, N.J., Griffin, W.L.	Garnets in a Polymict Xenolith from the Bultfontein Mine, South Africa: New Preliminary Geochemical and Textural Data.	968

Author	Title	Page No's
Wyatt, B.A., Sumpton, J.D.H., Shee, S.R., Smith, T.W	Kimberlites in The Forrest River Area. Kimberley Region, Western Australia.	971
Wyllie, P.J., Lee, W.J.	Kimberlites, carbonatites, peridotites and silicate-carbonate liquid immiscibility explained in parts of the system $\text{CaO}-(\text{Na}_2\text{O}+\text{K}_2\text{O})-(\text{MgO}+\text{FeO})-(\text{SiO}_2+\text{Al}_2\text{O}_3)-\text{CO}_2$	974
Yamashita, H., Arima, M., Ohtani, E.	Melting experiments of kimberlite compositions up to 9GPa: Determination of melt compositions using aggregates of diamond grains.	977
Yao, S., Griffin, W.L., O'Reilly, S.Y	Trace Elements in Chromites from Kimberlites and Related Rocks: Relation to Temperature and Mantle Composition.	980
Yaxley, G.M., Green, D.H.	Phase relations of carbonated eclogite under upper mantle PT conditions - implications for carbonatite petrogenesis.	983
Zack, T., Brumm, R	Ilmenite/liquid partition coefficients of 26 trace elements determined through ilmenite/clinopyroxene partitioning in garnet pyroxenites.	986
Zartman, R.E., Richardson, S.H., Gurney, J.J., Moore, R.O.	U-Th-Pb Ages of Megacrystic Zircon from the Monastery Kimberlite, Free State, South Africa.	989
Zhang, A., Griffin, W.L., Ryan, C.G., Andrew, A	Conditions of Diamond Formation beneath the Sino-Korean Craton: Paragenesis, Temperatures and the Isotopic Composition of Carbon.	992
Zhang, Y	Mechanical equilibria in inclusion-host systems	995
Zhao, D., Essene, E.J., Zhang, Y., Pell, J.A	Mantle Xenoliths from the Nikos Kimberlites on Somerset Island, and the Zulu Kimberlites on Brodeur Peninsula, Baffin Island, Canada.	998
Zhao, L., Zhang, P., Huang, X., Li, Y	Deep Mantle Fluids and Their Products in Kimberlites from China.	1001
Zheng, J.	Phanerozoic Evolution of the Subcontinental Lithospheric Mantle, Eastern North China Block: Mantle Xenolith Evidence.	1004
Zichella, V., de Gasparis, A.A., Pendock, N.E.	Mineral mapping with hyperspectral data. A case study over the Moses Rock Dyke and Mule Ear Diatreme (UTAH, USA)	1007
Zintchouk, N	New Data about Crusts of Weathering on Kimberlites of Diamondiferous Territories.	1009
Zintchouk, N.N., Boris, Y.I	Erosional Section of Kimberlite Bodies and the Scales of Placer Diamondiferousness.	1013
Zintchouk, N.N., Boris, Y.I., Stegnitsky, Y.B	Specific Features of Kimberlite Prospecting in Various Landscape-geological Conditions.	1017
Zintchouk, N.N., Dukardt, Y.A., Boris, Y.I	Specific Features of Zoning of Ancient Platforms' Territories According to the Degree of Perspectiveness of Diamondiferous Kimberlites' Intrusion.	1020
Zintchouk, N.N., Koptil, V.I., Boris, Y.I	Ancient Platforms' Diamond Typomorphism (on the example of Siberian Platform).	1024
Zintchouk, N.N., Zuev, V.M., Mitioukhine, S.I	Regional Zoning of Territories According to the Level of Primary Diamond Sources' Diamondiferousness.	1028
Zuev, V.M., Bezborodov, S.M., Chyerny, S.D., Yanygin, Y.T., Molchanov, Y.D	The Structures which Control the Location of Kimberlites of Middle-Markhinsky Region.	1031
Zuev, V.M., Serokurov, Y.N., Kalmykov, V.D	Assessment of Diamondiferous Perspectives of East-European Platform according to the Data of Sounding from Space.	1034
Zweistra, P., Jarvis, W., McGeorge, I.B.	The Geology of Micaceous Kimberlite Intrusives, Khutse, Botswana	1037

EXTENDED ABSTRACTS

